

# CLEANING LABORATORY EVALUATION SUMMARY

SCL #: 2009  
 DateRun: 04/24/2009  
 Experimenters: Jason Marshall, Junhee Cho  
 ClientType: Cleaner Manufacturer  
 ProjectNumber: Project #1  
 Substrates: Ceramics, Glass/Quartz, Stainless Steel  
 PartType: Coupon  
 Contaminants: Hucker's Soil, Food  
 Cleaning Methods: Low Pressure Spray  
 Analytical Methods: Gravimetric, Visual

Purpose: To evaluate performance of supplied automatic dishwashing formula for three soil types.

Experimental Procedure: A supplied product and rinse aid and a traditional automatic dishwashing formulation were tested to determine cleaning efficiencies for three soils from three substrates. The first two soils were taken from ASTM D3556. The third, Hucker's soil, was the all purpose soil the lab uses for janitorial testing.

Standard Food Soil Preparation: A mixture of 80 weight % of margarine and 20 weight % of powdered milk was prepared. The margarine was warmed until fluid and the powdered milk was mixed thoroughly.

Use of this optional soil makes the test more realistic by adding another difficult-to-remove component. Optional Food Soil Preparation: This mixture consists of 70 % margarine, 15 % powdered milk, and 15 % cooked cereal. The cooked cereal was separately prepared as follows: Add 45 g of cereal to 228 g of water, heat to boiling, and boil for 5 min. Dissolve 100 g of powdered milk in 500 g of water and stir this solution into the cooked cereal. Continue stirring as portions are removed to be combined with margarine and powdered milk that have been blended as in the previous soil preparation. This mixture was made up as required.

Hucker's Soil Formulation: Jif Creamy Peanut Butter 9.2%, Salted Butter 9.2%, Arrowhead Mills stone ground wheat flour 9.2%, Egg Yolk 9.2%, Evaporated milk 13.8%, Distilled water 45.8%, Printer's ink with boiled linseed oil 0.9%, Shaws saline solution 2.7%

Prew weighed ceramic (3), glass (3) and stainless steel (3) coupons (representing plates, glasses and silverware) were coated with each of the three soils and allowed to dry overnight. A second weight was recorded to determine the amount of soil added. All nine coupons were loaded into a VWR International Under-counter Glassware Washer (model 82020-922) and run on the light cycle (120 F wash, 140 F rinse). Total cleaning time lasted 2 hours. At the end of the cleaning, coupons were removed and weighed a final time to determine the amount of soil remaining. Percent efficiencies were calculated for each coupon cleaned. Observations were made for spotting or filming.

Results: When the ceramic coupons were weighed after cleaning, the final weights were greater than the initial weights. The weight changes may have been due to excess moisture being soaked into the uncoated side of the coupons, increasing the final weights. Visually, all the ceramic coupons for both cleaners were as clean as the glass and stainless steel coupons. Due to the weight change discrepancy, the gravimetric analysis of the ceramic coupons were not included in the overall efficiency calculations. The first table below lists the amount of soil added, the amount remaining and the efficiency for each coupon cleaned. The second table summarizes the cleaning performance of both product for each of three soils.

Cleaner	Initial wt	Final wt	% Removed
Alpha Chemical Service	0.1273	0.0017	98.66
Food soil #1	0.2438	0.0011	99.55
Glass	0.2409	0.0001	99.96
Alpha Chemical Service	0.2247	0.0003	99.87
Food soil #1	0.2914	-0.0001	100.03
Stainless	0.3269	0.0002	99.94
Alpha Chemical Service	1.5550	0.0010	99.94
Food soil #2	0.8081	0.0001	99.99
Glass	0.6418	0.0001	99.98
Alpha Chemical Service	1.0443	-0.0002	100.02
Food soil #2	1.0875	0.0006	99.94
Stainless	0.5575	0.0003	99.95

# CLEANING LABORATORY EVALUATION SUMMARY

Alpha Chemical Service	0.1015	0.0000	100.00
Hucker soil	0.0745	-0.0003	100.40
Glass	0.1208	0.0004	99.67
Alpha Chemical Service	0.0747	0.0004	99.46
Hucker soil	0.1112	-0.0004	100.36
Stainless	0.0770	0.0014	98.18
Cascade	0.1424	-0.0004	100.28
Food soil #1	0.1317	-0.0008	100.61
Glass	0.3055	-0.0002	100.07
Cascade	0.2059	0.0001	99.95
Food soil #1	0.1787	-0.0001	100.06
Stainless	0.1927	0.0000	100.00
Cascade	2.0406	-0.0002	100.01
Food soil #2	1.8063	-0.0002	100.01
Glass	1.2554	0.0000	100.00
Cascade	1.5470	0.0001	99.99
Food soil #2	0.5448	-0.0001	100.02
Stainless	0.9941	0.0001	99.99
Cascade	0.131	-0.0001	100.08
Hucker soil	0.2138	-0.0004	100.19
Glass	0.1341	-0.0005	100.37
Cascade	0.09	-0.0004	100.44
Hucker soil	0.1051	0.0004	99.62
Stainless	0.0887	0.0001	99.89

## Summary

	Food Soil 1	Food Soil 2	Hucker's Soil	Overall Average
Alpha Chemical Service	99.67	99.97	99.68	99.77
Cascade	100.16	100	100.1	100.09
Ceramic observations				
Alpha Chemical Service	Pass	Pass	Pass	
Cascade	Pass	Pass	Pass	

Summary:

<b>Substrates:</b>	Ceramics, Glass/Quartz, Stainless Steel				
<b>Contaminants:</b>	Hucker's Soil, Food				
<b>Company Name:</b>	<b>Product Name:</b>	<b>Conc.:</b>	<b>Efficiency:</b>	<b>Effective:</b>	<b>Observations:</b>
Procter & Gamble	Cascade Complete (Dawn)		100.09	<input checked="" type="checkbox"/>	
Alpha Chemical Services	Alpha Automatic Dish Detergent		99.77	<input checked="" type="checkbox"/>	

Conclusion:

The supplied product worked as well as the traditional cleaning product removing close to 100% of each of the three soils.